

## Paediatric acquired demyelinating syndromes : incidence, clinical and magnetic resonance imaging features

Absoud, M; Lim, MJ; Chong, WK; De Goede, CG; Foster, K; Gunny, R; Hemingway, C; Jardine, Phillip; Kneen, R; Likeman, M; Nischal, KK; Pike, MG; Sibtain, NA; Whitehouse, WP; Cummins, Carole; Wassmer, E

DOI:

[10.1177/1352458512445944](https://doi.org/10.1177/1352458512445944)

License:

Creative Commons: Attribution-NonCommercial (CC BY-NC)

*Document Version*

Publisher's PDF, also known as Version of record

*Citation for published version (Harvard):*

Absoud, M, Lim, MJ, Chong, WK, De Goede, CG, Foster, K, Gunny, R, Hemingway, C, Jardine, P, Kneen, R, Likeman, M, Nischal, KK, Pike, MG, Sibtain, NA, Whitehouse, WP, Cummins, C & Wassmer, E 2013, 'Paediatric acquired demyelinating syndromes : incidence, clinical and magnetic resonance imaging features', *Multiple Sclerosis*, vol. 19, no. 1, pp. 76-86. <https://doi.org/10.1177/1352458512445944>

[Link to publication on Research at Birmingham portal](#)

### **Publisher Rights Statement:**

Eligibility for repository : checked 24/02/2014

### **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### **Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

# Multiple Sclerosis Journal

<http://msj.sagepub.com/>

---

## **Paediatric acquired demyelinating syndromes: incidence, clinical and magnetic resonance imaging features**

Michael Absoud, Ming J Lim, Wui K Chong, Christian G De Goede, Katharine Foster, Roxana Gunny, Cheryl Hemingway, Philip E Jardine, Rachel Kneen, Marcus Likeman, Ken K Nischal, Michael G Pike, Naomi A Sibtain, William P Whitehouse, Carole Cummins, Evangeline Wassmer and The UK and Ireland Childhood CNS Inflammatory Demyelination Working Group

*Mult Scler* 2013 19: 76 originally published online 19 April 2012

DOI: 10.1177/1352458512445944

The online version of this article can be found at:

<http://msj.sagepub.com/content/19/1/76>

---

Published by:



<http://www.sagepublications.com>

**Additional services and information for *Multiple Sclerosis Journal* can be found at:**

**Open Access:** Immediate free access via SAGE Choice

**Email Alerts:** <http://msj.sagepub.com/cgi/alerts>

**Subscriptions:** <http://msj.sagepub.com/subscriptions>

**Reprints:** <http://www.sagepub.com/journalsReprints.nav>

**Permissions:** <http://www.sagepub.com/journalsPermissions.nav>


>> [Version of Record](#) - Jan 9, 2013

[OnlineFirst Version of Record](#) - Apr 19, 2012

[OnlineFirst Version of Record](#) - Apr 19, 2012

[What is This?](#)

# Paediatric acquired demyelinating syndromes: incidence, clinical and magnetic resonance imaging features

*Multiple Sclerosis Journal*  
19(1) 76–86  
© The Author(s) 2012  
Reprints and permissions:  
sagepub.co.uk/journalsPermissions.nav  
DOI: 10.1177/1352458512445944  
msj.sagepub.com  


Michael Absoud<sup>1</sup>, Ming J Lim<sup>2</sup>, Wui K Chong<sup>3</sup>,  
Christian G De Goede<sup>4</sup>, Katharine Foster<sup>5</sup>, Roxana Gunny<sup>3</sup>,  
Cheryl Hemingway<sup>6</sup>, Philip E Jardine<sup>7</sup>, Rachel Kneen<sup>8</sup>,  
Marcus Likeman<sup>9</sup>, Ken K Nischal<sup>10</sup>, Michael G Pike<sup>11</sup>,  
Naomi A Sibtain<sup>12</sup>, William P Whitehouse<sup>13</sup>, Carole Cummins<sup>1\*</sup>  
and Evangeline Wassmer<sup>14\*</sup> for The UK and Ireland Childhood  
CNS Inflammatory Demyelination Working Group

## Abstract

**Objective:** Changing trends in multiple sclerosis (MS) epidemiology may first be apparent in the childhood population affected with first onset acquired demyelinating syndromes (ADSs). We aimed to determine the incidence, clinical, investigative and magnetic resonance imaging (MRI) features of childhood central nervous system ADSs in the British Isles for the first time.

**Methods:** We conducted a population active surveillance study. All paediatricians, and ophthalmologists ( $n = 4095$ ) were sent monthly reporting cards (September 2009–September 2010). International Paediatric MS Study Group 2007 definitions and McDonald 2010 MS imaging criteria were used for acute disseminated encephalomyelitis (ADEM), clinically isolated syndrome (CIS) and neuromyelitis optica (NMO). Clinicians completed a standard questionnaire and provided an MRI copy for review.

**Results:** Card return rates were 90%, with information available for 200/222 positive notifications (90%). After exclusion of cases, 125 remained (age range 1.3–15.9), with CIS in 66.4%, ADEM in 32.0% and NMO in 1.6%. The female-to-male ratio in children older than 10 years ( $n = 63$ ) was 1.52:1 ( $p = 0.045$ ). The incidence of first onset ADS in children aged 1–15 years old was 9.83 per million children per year (95% confidence interval [CI] 8.18–11.71). A trend towards higher incidence rates of ADS in children of South Asian and Black ethnicity was observed compared with White children. Importantly, a number of MRI characteristics distinguished ADEM from CIS cases. Of CIS cases with contrast imaging, 26% fulfilled McDonald 2010 MS diagnostic criteria.

<sup>1</sup>School of Health and Population Sciences, University of Birmingham, Birmingham, UK.

<sup>2</sup>Department of Neurology, The Evelina Children's Hospital at Guy's and St Thomas' NHS Trust, London, UK.

<sup>3</sup>Department of Neuroradiology, Great Ormond Street Hospital for Children, London, UK.

<sup>4</sup>Department of Paediatric Neurology, Royal Preston Hospital, Lancashire, UK.

<sup>5</sup>Department of Neuroradiology, Birmingham Children's Hospital, Birmingham, UK.

<sup>6</sup>Department of Neurology, Great Ormond Street Hospital for Children, London, UK.

<sup>7</sup>Department of Neurology, Bristol Royal Hospital for Children, Bristol, UK.

<sup>8</sup>Department of Neurology, Alder Hey Children's NHS Foundation Trust, Liverpool, UK.

<sup>9</sup>Department of Neuroradiology, Bristol Royal Hospital for Children, Bristol, UK.

<sup>10</sup>Department of Ophthalmology, Great Ormond Street Hospital for Children, London, UK.

<sup>11</sup>Department of Neurology, Oxford Children's Hospital, Oxford, UK.

<sup>12</sup>Department of Neuroradiology, King's College Hospital NHS Trust, London, UK.

<sup>13</sup>School of Clinical Sciences, University of Nottingham, Nottingham, UK.

<sup>14</sup>Department of Neurology, Birmingham Children's Hospital, Birmingham, UK.

\*Joint last authors

## Corresponding author:

Dr Michael Absoud, School of Health and Population Sciences, University of Birmingham, Vincent Drive, Birmingham B15 2TT, UK  
Email: michaelabsoud@childdemyelination.org.uk

**Conclusions:** We report the highest surveillance incidence rates of childhood ADS. Paediatric MS diagnosis at first ADS presentation has implications for clinical practice and clinical trial design.

### Keywords

pediatric multiple sclerosis, paediatric multiple sclerosis, acute disseminated encephalomyelitis, demyelination, demyelinating, pediatric, paediatric, childhood, adolescent, epidemiology, incidence, magnetic resonance imaging

Date received: 27th February 2012; revised: 21st March 2012; accepted: 27th March 2012

## Introduction

Multiple sclerosis (MS) is a chronic disabling and costly neurodegenerative disorder. Incidence of MS appears to be increasing worldwide with a rising female-to-male sex ratio,<sup>1,2</sup> and with approximately 5% of adults who develop MS having first symptoms in childhood.<sup>3,4</sup> Some epidemiological evidence suggests early environmental exposures may influence MS risk.<sup>5,6</sup> Hence, any epidemiological change in MS may be evident first in a paediatric population.

Diagnosis of MS in children, as in adults, is first heralded by onset of a clinically isolated syndrome (CIS), or more rarely acute disseminated encephalomyelitis (ADEM). Children diagnosed with relapsing–remitting MS present with one of the central nervous system (CNS) acquired demyelinating syndromes (ADS) followed by a second event usually within 2 years.<sup>7</sup> Childhood CIS presentations with an abnormal magnetic resonance imaging (MRI) of the brain (i.e. more than one high T2 signal) compared with ADEM are more likely to have relapsing disease.<sup>8–12</sup> Recently, the new MRI McDonald 2010 MS criteria have been recommended for use in children presenting with a CIS,<sup>13</sup> potentially allowing for MS diagnosis at first presentation of a CIS.<sup>14</sup> There is no biomarker that differentiates between monophasic ADEM, monophasic CIS, or MS, hence a combination of clinical, cerebrospinal fluid (CSF) positive oligoclonal bands and MRI features is suggested.<sup>12, 15</sup> MRI differences, in particular between ADEM and polyfocal CIS, are also not well recognized.<sup>16</sup> The differential diagnosis of ADS is complex and wide, especially so in childhood.<sup>17</sup>

The International Paediatric Multiple Sclerosis Study Group (IPMSSG) have published consensus definitions (2007) of paediatric CNS ADS and MS to facilitate uniformity in clinical practice and future epidemiological research.<sup>4</sup> Two previous surveillance studies in Canada (2004–2007) and Germany (1997–1999) revealed an incidence ranging from 3.7<sup>18</sup> to 9.0 per million children per year.<sup>19</sup> The incidence of childhood ADS remains largely unknown worldwide, and as the incidence of these disorders based on the epidemiology of MS in adults is likely to differ geographically, estimates are needed from multiple geographical regions.

We report a multinational multisource<sup>20</sup> prospective active surveillance study in the British Isles (latitudes

50–59°N) to ascertain incidence, clinical, investigative and MRI features of childhood (age <16 years) ADS using panel review to apply IPMSSG consensus definitions (including the incorporated McDonald MRI 2001 and Barkhof space diagnostic criteria).<sup>4, 21</sup> We also apply the newly revised 2010 McDonald space and time diagnostic criteria at first ADS presentation;<sup>13</sup> and the recent Verhey MRI prognostic MS criteria.<sup>22</sup>

## Methods

### Study design

We carried out a prospective active surveillance study of first episode ADS in children aged between 1 and 15 years 11 months of age, in the British Isles. We used multisource ascertainment via two well-established surveillance units in order to establish incidence of new cases.

### *The British Paediatric Surveillance Unit (BPSU), the primary source of data*

The British Paediatric Surveillance Unit (BPSU) facilitates epidemiological surveillance of childhood diseases of public health importance in the UK and Republic of Ireland. The BPSU, launched in 1986, has helped initiate over 80 studies with a card return rate consistently higher than 93%.

### *The British Ophthalmological Surveillance Unit (BOSU), the alternative source of data for optic neuritis cases*

The British Ophthalmological Surveillance Unit (BOSU) enables ophthalmologists to participate in the nationwide surveillance of uncommon ophthalmological conditions. The BOSU card return rate is consistently higher than 75%.

### *Case ascertainment via active surveillance*

One month before commencing the study, a protocol card and letter was sent to all clinicians informing them of the study. Monthly notification cards (mean 4095/month) were sent by the BPSU (mean 2945/month) to all registered certified consultant paediatricians and paediatric neurologists

**Table 1.** Summarized inclusion definitions for CNS acquired demyelinating syndromes.

Condition	Definition
Acute disseminated encephalomyelitis (ADEM)	(1) A polysymptomatic clinical event with acute/subacute onset that must include encephalopathy (behavioural change or altered consciousness). (2) MRI brain shows multifocal lesions.
Clinically isolated syndrome (CIS)	A first acute-clinical episode of CNS symptoms which may either be monofocal or multifocal, but does not include encephalopathy (except in brainstem syndromes). The MRI will show white matter demyelination. These include: <ol style="list-style-type: none"> <li>1. Transverse myelitis (TM): weakness and/or numbness of both legs +/- arms, usually with maximal deficits 1 week after symptom onset supported by demyelination on MRI spine.</li> <li>2. Optic neuritis: Acute or subacute loss of vision and <math>\geq 1</math> of: relative afferent pupillary defect (unilateral cases), visual field deficit or scotoma, impaired colour vision, optic disc oedema, or abnormal visual evoked potentials. MRI is not necessary for diagnosis.</li> <li>3. Other CIS: Brainstem, cerebellar, and/or hemispheric dysfunction, supported by demyelination on MRI.</li> </ol>
Neuromyelitis optica (NMO)	Must have: i. Optic neuritis and ii. Acute myelitis. Must have: iii. Spinal MRI lesion extends over three or more segments or iv. Aquaporin-4 antibody testing is positive. The brain MRI may be abnormal but must not meet Multiple Sclerosis MRI diagnosis criteria.
Exclusion criteria	<ol style="list-style-type: none"> <li>1. Leukodystrophies (e.g. metachromatic leukodystrophy, adrenoleukodystrophy) or mitochondrial disease.</li> <li>2. Proven CNS infection (e.g. viral encephalitis, bacterial meningitis, herpes simplex encephalitis, Lyme disease, HIV).</li> <li>3. Radiation/chemotherapy associated white matter damage.</li> <li>4. Condition fulfilling criteria for CNS connective tissue disease e.g. lupus, vasculitis. The sole presence of antibodies associated with CNS connective tissue or autoimmune diseases was not sufficient for exclusion.</li> </ol>

and BOSU (mean 1150/month) to all registered ophthalmologists (between September 2009 and September 2010). In the British Isles, these clinicians are likely to see all cases of childhood ADS. Clinicians returned the card to the surveillance units notifying any cases or confirming 'nothing to report'. Upon receipt of a positive notification the surveillance units provided the investigating team with a BPSU/BOSU case number and clinician contact details. The study team contacted the clinician directly, sending a detailed data collection form and a blank CD for an anonymized MRI. As more than one clinician might notify the same patient seen in different settings, minimal identifiers (National Health Service [NHS] number, district post-code, sex and date of birth) were used for record linkage and to exclude duplicates with a high level of certainty (<http://www.rcpch.ac.uk/bpsu>).

### Initial surveillance reports

Using the IPMSSG consensus definitions for classification of paediatric ADS,<sup>4</sup> we asked clinicians to report children experiencing clinical neurological events consistent with a first episode ADS and confirmed with white matter changes (except in optic neuritis) on MRI (Table 1).

### Data collection

A condition specific targeted neurological history was collected at the time of first demyelinating event. Data collected included: basic demographic data; demyelinating symptoms and signs; CSF findings, serological and

microbiological results; timing of MRI being carried out and whether abnormal; and treatment given.

### MRI imaging review

MRI scans were reviewed blinded to clinical features by MA and one of five neuroradiologists (KC, KF, RG, ML, NS). A standardized proforma was completed utilizing previously described nomenclature (supplementary material appendix 1).<sup>10, 23, 24</sup> McDonald 2001 dissemination in space criteria (as recommended by the current IPMSSG criteria),<sup>4, 21</sup> McDonald 2010 dissemination in space and time (if a gadolinium-enhanced scan was available) criteria and Verhey prognostic criteria were classified by MA according to the MRI variables and associated panel review.<sup>13</sup> Features of ADEM and CIS scans were compared. Although a single MRI protocol was not feasible as this was a national population based study collecting data from established clinical practice from all centres, all MRI brain scans were performed on 1.5 T scanners and had a minimum of a T1-weighted and a T2-weighted or fluid-attenuated inversion recovery (FLAIR) sequence in two different planes. Additional sequences included gadolinium-enhanced T1, diffusion weighted and spine images. MRI slice thickness varied from 3–5 mm.

### Panel classification and agreement for inclusion

The co-investigators met quarterly as a panel to review reported cases (at least four members present at each meeting). The panel consisted of paediatric neurologists ( $n = 8$ ;

**Table 2.** Incidence of childhood CNS acquired demyelinating syndromes by country, sensitivity analysis and ethnic group in England and Wales.

CONFIRMED CASES						
Region	Number	Percentage	Population 2010 1–15 years (per thousands) or %	Incidence per annum per million	95% CI lower	95% CI upper
England, Wales and Channel Islands	117	93.6	9636	11.2	9.27	13.4
Scotland	3	2.4	853	3.25	0.65	9.49
Ireland and Northern Ireland	5	4.0	1254	3.68	1.19	8.59
Total	125	100.0	11743	9.83	8.18	11.71
<b>Sensitivity Analysis Assuming All Reports With Missing Information Are Cases</b>						
England, Wales and Channel Islands	129	87.8	9636	12.36	10.32	12.68
Scotland	9	6.1	853	9.74	4.44	18.5
Ireland and Northern Ireland	9	6.1	1254	6.63	3.0	12.6
Total	147	100.0	11743	11.6	9.76	13.6
<b>Incidence By Ethnicity In England And Wales</b>						
White	93	80.17%	83.93%	10.62	8.57	13.00
South Asian: Indian; Pakistani; Bangladesh; other South Asian background	11	9.48%	7.30%	14.43	7.20	25.83
Black	7	6.03%	3.31%	20.25	8.12	41.73
Chinese, mixed and other	5	4.31%	5.46%	8.78	2.82	20.27
Total	116	116	100%			

CDG, CH, EW, MJL, MP, PJ, RK, WPW) with expertise in ADS, and final classification as per IPMSSG clinical criteria was arrived at by consensus. MRI images (with neuroradiologist scoring) were used to confirm the presence of lesions consistent with inflammatory demyelination, to aid in the exclusion of cases and to sub-classify CIS cases with asymptomatic MRI lesions (supplementary material appendix 1).<sup>17</sup>

### Statistical analysis

Descriptive statistics were used to summarize the key components of the dataset. Estimates of national incidence with confidence intervals (Byar's approximation of the exact Poisson) for the 13-month study was annualized using mid-2010 UK and 2010 Republic of Ireland population estimates.<sup>25,26</sup> Mid-2009 population estimates by ethnicity for England and Wales were used to evaluate rates of childhood ADS by ethnicity (Table 2). A sensitivity analysis was conducted on a worst-case scenario that all reported cases with missing clinical histories were actually cases of ADS to assess the potential impact on estimated incidence of ADS. We used a two source case capture–recapture analysis to estimate missed cases of optic neuritis on the assumption of independence,<sup>27</sup> but did not carry out capture–recapture analysis for other ADS, as referral networks mean that the assumption of independence was violated. The England 2010 index of multiple deprivation was used to derive the percentage of cases in England living in the 20% most deprived districts.<sup>28</sup> Parametric or non-parametric statistical tests (Kruskal–Wallis tests) were used for continuous distributions as appropriate given normality and  $\chi^2$  or Fisher's exact tests for nominal data. Agreement

between reporters and the panel was measured using the Kappa statistic. Statistical analysis was performed using PASW Statistics for Windows version 17.0 (© SPSS, Inc., 2009, Chicago, IL, www.spss.com) and the openepi software (version 2.3.1, www.openepi.com).<sup>29</sup>

### Approvals


The study was approved by the BPSU and BOSU executive committees. Data handling was restricted to those with direct involvement in the project. To comply with collection of minimal identifier information regulations without patient consent as in Section 251 of the NHS Act 2006, the study has approval from the National Information Governance Board's Ethics & Confidentiality Committee (ECC/BPSU 4-03[FT1]/2009) and the UK Multicentre Research Ethics Committee (09/H1202/92).

### Results

#### Card return rate, expert review and record linkage outcome

Card return rate was 94% from the BPSU and 78% from the BOSU (total 47,910 from 53,235 cards returned including negative reports). There were 222 positive notifications (197 BPSU and 25 BOSU) with insufficient information for review of 22 notifications (6 Scotland, 12 England and Wales, and 4 Ireland and Northern Ireland). In the remaining 200 (90%), 41 duplicate notifications were identified. Cases where the event was outside the inclusion dates, September 2009 to September 2010 ( $n = 16$ ) were excluded. A further 18 cases were excluded by the panel (Figure 1). A





British Paediatric Surveillance Unit (BPSU); British Ophthalmological Surveillance Unit (BOSU);  
N-Methyl-D-aspartate receptor (NMDAR).

**Figure 1.** Study flow chart showing panel review and record linkage outcome.

total of 124/125 cases remaining had MRI scans (one isolated optic neuritis had no MRI scan, four transverse myelitis cases had MRI spine only).

### *Incidence of childhood CNS inflammatory demyelination in the British Isles*

The minimum incidence of first onset ADS in children aged 1–15 years old in the British Isles is 9.83 per million

children per year (95% confidence interval [CI] 8.18–11.71). Incidence was highest in England and Wales (Table 2). A sensitivity analysis showed no significant difference in incidence rates within countries in the British Isles.

### *Demographic characteristics*

Median age was 10.0 years (range 1.3–15.9; mean 9.74 years, SD 4.34) and 51.2% ( $n = 64$ ) of cases were female

**Table 3.** Clinical, demographic, and phenotypic expert panel classification of the childhood CNS acquired demyelinating syndromes.

		CIS (n = 83)				
		ADEM (n = 40)	ON (n = 31)	TM (n = 26)	Other CIS (n = 26)	NMO (n = 2)
<b>Age at presentation: median (IQR)</b>		5.3 (3.6-7.0)	11.8 (9.0-13.9)	12.6 (9.3-14.0)	14.0 (9.5-14.5)	6.4 and 14.8 years
<b>Sex (n = 125)</b>						
	<b>Male: Female</b>					
	<b>(% female) 61: 64</b>					
	<b>(51%)</b>	24:16 (40%)	15:16 (52%)	11:15 (58%)	11:15 (58%)	0:2 (100%)
<b>Ethnicity (n = 124/125)</b>				37:46 (55%)		
	<b>White (n = 101)</b>	36 (90%)	21 (68%)	22 (85%)	21 (81%)	1 (50%)
	<b>Asian (n = 11)</b>	3	4	2	1	1
	<b>Black (n = 7)</b>	1	3	1	2	0
	<b>Chinese, mixed, other (n = 5)</b>	0	3	1	1	0
<b>Clinical:</b>						
Monofocal	<b>MRI:</b>					
	≥ 1 asymptomatic lesion		7 (23%)	9 (35%)	4 (15%)	
Multifocal	no asymptomatic		24	17	3 (12%)	
	≥ 1 asymptomatic		0	0	16 (62%)	
Radiologically isolated	no asymptomatic		0	0	2 (8%)	
			n/a	n/a	1	
<b>Spinal lesion(s) present: n (% ≥ 3 segments)</b>		8/12 (100%)	1/8 (0)	26/26 (62%)	4/9 (75%)	2/2 (100%)
<b>Death</b>		1/40 (2.5%)	0	0	0	0
<b>Seizures</b>		8/40 (20%)	0	0	0	0
<b>Optic neuritis: Unilateral: bilateral (% bilateral)</b>		0:3 (100%)	22:9 (29%)	0:0	0:1	1:1 (50%)
<b>Cerebellar signs: n (%)</b>		18 (45%)	0	0	11 (42%)	0
<b>Brain stem signs: n (%)</b>		11 (28%)	0	1	13 (50%)	0
<b>Pyramidal signs: n (%)</b>		24 (60%)	0	0	10 (38%)	0
<b>Intensive care unit admissions; n = 12 (10%)</b>		8 (20%)	0/31 (0%)	3/26 (12%)	1/26 (4%)	0/2
<b>Plasma exchange; n = 4 (3%)</b>		2 (5%)	0/31	2/26 (8%)	0/26	0/2
<b>Intravenous corticosteroids</b>		37/40 (93%)	23/31 (74%)	25/26 (96%)	18/26 (69%)	2/2 (100%)
<b>Oral steroids</b>		27/40 (68%)	21/31 (67%)	16/26 (62%)	11/26 (42%)	2/2 (100%)
<b>Intravenous immunoglobulin</b>		6/40 (15%)	0	7/26 (27%)	1/26 (4%)	0

ADEM, acute disseminated encephalomyelitis; ON, optic neuritis; TM, transverse myelitis; CIS, clinically isolated syndrome; NMO, neuromyelitis optica; MRI, magnetic resonance imaging; IQR, interquartile range.


(Table 3). The median age for males was 8.9 years (interquartile range [IQR] 5.0–13.5) and for females was 11.4 years (IQR 6.8–14.4;  $p = 0.046$ ). The female-to-male ratio in children younger than 10 years ( $n = 62$ ) was 0.72:1 and in older than 10 years ( $n = 63$ ) was 1.52:1. White children comprised 81% of the cohort. Higher incidence rates of ADS in children of South Asian and Black ethnicity in England and Wales compared with White children were observed, although not reaching statistical significance (Table 2). A total of 34% of the cases in England lived in the most 20% deprived districts. Children living in the most 20% deprived areas were more likely to be males compared with female (23:13) when compared with those living in the 80% least deprived areas (30:40,  $p = 0.040$ ). They were also most likely to be of non-White ethnicity (12:24 versus

11:58,  $p = 0.041$ ). Age and ADS presentation did not influence those living in the most 20% deprived areas.

### Results of panel review and clinical classification

Childhood ADS were classified as CIS in 66.4%, ADEM in 32.0% and neuromyelitis optic (NMO) in 1.6%. Of the CIS cases, optic neuritis (ON) represented 37.4%, transverse myelitis (TM) 31.3% and other CIS 31.3% (Table 3). The majority (70%) of the other CIS cases were polysymptomatic. Children with ADEM were younger than cases presenting with CIS ( $p < 0.001$ ) (Figure 2). The risk ratio for CIS in females compared with males was 1.46 (95% CI 1.17–1.83). The panel reclassified a total of 15 out of the





**Figure 2.** Childhood acquired demyelinating syndromes by age (younger and greater than 10 years old) and sex.

125 cases (12%): ADEM to CIS ( $n = 11$ ) mainly due to lack of encephalopathy; Other CIS to TM ( $n = 3$ ); TM to other CIS ( $n = 1$ ). The overall level of agreement between reporters and the panel was good (Kappa 0.838 [95% CI 0.761–0.914]).

## Deaths

Clinicians reported three deaths in the initial hospital admission, one in a child with ADEM and two in cases which proved not to have ADS (post-transplant lymphoproliferative disorder and hypereosinophilic syndrome).

## Capture–recapture analysis

Capture–recapture analysis for ON cases ( $n = 31$ ) assuming that BPSU and BOSU were independent sources of cases estimated the total population to be 47 (95% CI 30–55).

## MRI features

More CIS cases fulfilled MRI McDonald 2010 dissemination in space criteria (42%) compared with the 2001 criteria (28%; Table 4). Of the CIS cases which had contrast-enhanced scans ( $n = 38$ ), 10 cases (26%) fulfilled McDonald 2010 space and time criteria for MS diagnosis. Verhey MRI prognostic criteria for identifying childhood ADS at high risk of MS (presence of one T1 hypointense and one periventricular lesion) were fulfilled in 11/78 (14%) CIS cases compared with 1/40 (2.5%) ADEM cases. A total of 19/78 (24%) CIS cases fulfilled either Verhey prognostic criteria or McDonald 2010 MS space and time criteria for MS (8/10 cases which fulfilling McDonald 2010 space and time criteria did not fulfil Verhey criteria, whilst 3/5 fulfilling Verhey criteria did not fulfil McDonald criteria).

ADEM cases compared with CIS cases with abnormal MRI (supplementary material appendix 2) were more likely to have more deep grey nuclei ( $p < 0.001$ ), large white matter ( $p < 0.001$ ) and cortical grey matter ( $p = 0.001$ ) high T2 signal lesions. CIS scans were more likely to have more periventricular ( $p < 0.001$ ), deep white matter ( $p = 0.003$ ), corpus callosum ( $p = 0.001$ ), well-defined and discrete high T2 signal lesions ( $p < 0.001$ ) and black holes ( $p = 0.008$ ). The number of total high T2 signal, enhancing, infratentorial, juxtacortical and spinal lesions did not differentiate between CIS and ADEM patients. Of the ON and TM cases (monofocal CIS) 23% and 35% had one or more high T2 signal brain lesion respectively (Table 3).

## Investigative features

Children with ADEM more commonly had CSF lymphocytosis ( $p = 0.005$ ) and less commonly CSF positive oligoclonal bands ( $p = 0.011$ ) compared with those with CIS (Table 5). One ADEM case was reported as having the common missense mutation (c.1880CT, p.Thr585Met) of the Ran Binding Protein 2 (RANBP2) gene previously reported in recurrent or familial

**Table 4.** MRI criteria fulfilled.

	Panel classification		
	CIS ( $n = 83$ )		
	ON ( $n = 31$ )	TM ( $n = 26$ )	Other CIS ( $n = 26$ )
<b>McDonald 2001 space criteria</b>	4/30	5/22	13/26
<b>McDonald 2010 space criteria</b>	5/30	8/22	20/26
<b>McDonald 2010 time criteria (gadolinium given scans)</b>	0/12	3/11	7/15
<b>Meets McDonald MRI criteria for MS on initial presentation (dissemination in space and time)</b>		10/38 (26%)	
<b>Verhey or McDonald 2010 MS criteria present</b>		19/78 (24%)	

MRI, magnetic resonance imaging; MS, multiple sclerosis; ON, optic neuritis; TM, transverse myelitis; CIS, clinically isolated syndrome.

**Table 5.** Investigative features.

	Expert panel classification				
	ADEM ( <i>n</i> = 40)	CIS ( <i>n</i> = 83)			NMO ( <i>n</i> = 2)
		ON ( <i>n</i> = 31)	TM ( <i>n</i> = 26)	Other CIS ( <i>n</i> = 26)	
<b>CSF oligoclonal bands</b>	2/20 (10%)	6/17 (35%)	6/19 (32%)	10/15 (67%)	0/1 (0%)
<b>CSF WBC <math>\geq 5 \times 10^6/l</math></b>	21/36 (58%)	3/17 (18%)	8/20 (40%)	8/21 (38%)	0/2 (0%)
<b>CSF protein median (IQR)</b>	0.38 (0.25–0.58) <i>n</i> = 28	0.26 (0.23–0.35) <i>n</i> = 17	0.36 (0.26–0.52) <i>n</i> = 19	0.29 (0.21–0.36) <i>n</i> = 17	0.29 <i>n</i> = 2
<b>Genetic tests reported</b>	1 case RANBP2 positive	n/a	n/a	n/a	n/a

ADEM, acute disseminated encephalomyelitis; ON, optic neuritis; TM, transverse myelitis; CIS, clinically isolated syndrome; NMO, neuromyelitis optica; CSF, cerebrospinal fluid RANBP2, Ran Binding Protein 2 gene; IQR, interquartile range; WBC, white blood cell.

acute necrotising encephalopathy.<sup>30</sup> Of those tested for the aquaporin-4 antibody (*n* = 52), none of the ADEM (*n* = 5), TM (*n* = 20), and other CIS (*n* = 7) cases tested positive while one child with ON and both NMO patients tested positive.

### Treatment

The majority (89%) received either intravenous corticosteroids (84%) and/or oral prednisolone in the first instance. A higher proportion of children with ADEM were required intensive care unit (ICU) admission, and four children (two ADEM and two TM) received plasma exchange. Intravenous immunoglobulin was administered to 11% of the cohort (Table 3).

### Discussion

The minimum incidence of first onset childhood ADS in the British Isles is 9.83 per million (95% CI 8.18–11.71) children per year aged 1–15 years. This is the highest reported incidence of childhood ADS from a prospective childhood surveillance study. To date, there have been two surveillance studies studying ADS. A Canadian surveillance study (2004–2007) estimated the incidence of childhood ADS to be 9.0 per million people aged less than 18 years old,<sup>19</sup> with 22% classified as ADEM and 78% as CIS. Although this study did not have MRI review and was conducted prior to the IPMSSG consensus definitions, it is unlikely that these factors would have had a major influence on classification. An earlier surveillance study from Germany (1997–1999) estimated the incidence of MS in children <16 years as 3.0 per million children, and that of ADEM as only 0.7 per year per million children aged less than 16 years old.<sup>18</sup> This study asked clinicians to report new cases of definite or suspected MS or ADEM cases, and did not include other ADS. Diagnosis of MS and ADS depends on clinical definitions and MRI criteria used, and hence it may be that these ascertainment estimates would change with current definitions and criteria (supplementary material appendix 3).

The mean age of onset (9.7 years) was similar to the Canadian incidence study (mean 10.5 years) and a French<sup>9</sup> (children <16 years, 1985–2001) first episode ADS retrospective cohort study (mean 9.9 years), but older than a Dutch<sup>10</sup> (children <16 years, 1990–2010) retrospective cohort study (8.5 years). A recent retrospective study from the USA (*n* = 85) showed the incidence of childhood ADS and MS to be higher in Black children compared with White and Hispanic children.<sup>31</sup> While non-White ethnic groups were over-represented in our study (*n* = 125), this did not reach statistical significance. Although there is limited power to compare subgroups in the study, this still contrasts with the adult MS population where white ethnic groups are over-represented. Adult and paediatric<sup>7,14,32</sup> MS studies have shown that adolescent females are over-represented compared with males with ratios of up to 3:1. An adult CIS study has also shown a higher risk ratio of females developing CIS,<sup>33</sup> the first presentation of MS. Our study shows that in childhood first onset ADS, sex ratios also varied with age, with more females in the older age groups (F:M ratio of 1.52:1 in children over 10 years). Children older than 10 years more often had CIS compared with those less than 10 years of age who presented more often with ADEM. A Japanese retrospective multicenter study showed the male-to-female ratio of ADEM to be 2.3:1, and that ADEM was three times more common than MS.<sup>34</sup> In addition previous UK epidemiological adult studies have shown that MS hospital admissions, incidence and prevalence were higher in those living in least deprived areas.<sup>35,36</sup> Our study showed that children with ADS living in the most 20% deprived areas (in England) were more likely to be males and of non-White ethnicity. These observations regarding the geographical deprivation indices and demographics of childhood ADS (over-representation of older females with CIS, and children with Black and South Asian ethnicity) add further support to the hypothesis that environmental influences may indeed be acting as early as in childhood.

This study is not without limitations. As with other epidemiological studies, some cases may not have been reported or have had sub-clinical presentations, and it is plausible that children in our study who met McDonald

2010 criteria for MS at first clinical presentation may have had previous symptoms not identified as ADS. This was addressed as far as was possible by using clear consensus case definitions and multiple sources of case ascertainment. A capture–recapture analysis estimated that a possible 16 ON cases were missed out of a potential 47, although this relies on assumptions of complete independence that may not hold in practice.

Our study shows that childhood CIS occurs more often than ADEM (ratio 2.1:1) and also confirms a previous report that ADEM can be a severe disease with one death and one in five requiring intensive care admission.<sup>37</sup> The proportion of children with ADEM in our cohort (32%) was slightly higher than French<sup>9</sup> (29%), Canadian (22%), and Dutch<sup>10</sup> cohorts (termed polyfocal ADS with encephalopathy, not ADEM).

We found that 43% of CIS patients had one or more asymptomatic brain lesion. Previous studies have shown that children with CIS with abnormal brain scans are at a higher risk of relapse than those without asymptomatic brain lesions.<sup>11,12</sup> Also it is of interest that a number of MRI measures significantly helped differentiate ADEM from CIS cases. For example, ADEM cases compared with CIS had more deep grey nuclei and cortical grey matter high T2 signal lesions. CIS cases had more periventricular, deep white matter, corpus callosum high T2 signal lesions and black holes.

CSF lymphocytosis and negative oligoclonal bands were associated with an ADEM diagnosis, in contrast to a previous study.<sup>16</sup> It is of interest that one case was reported as positive for the common mutation in the RANBP2 gene, which has been recently described in familial or recurrent acute necrotizing encephalopathy (ANE).<sup>30</sup> With the increasing recognition of the overlap between the MRI and clinical features of ADEM and ANE, and further reports of other genetically predetermined encephalopathies,<sup>38</sup> this potential novel group of genetic determinant(s) of encephalopathy in demyelination requires exploration.

A total of 26% of CIS cases who had gadolinium given scans ( $n = 38$ ) fulfilled McDonald 2010 MRI criteria for MS at onset (space and time), hence a minimum 12% of CIS patients met McDonald 2010 MS diagnostic criteria at diagnosis. In addition 24% of CIS cases fulfilled either McDonald 2010 or Verhey criteria which are thought to indicate higher risk of MS, although both the McDonald criteria and the Verhey model require further validation in prospective multinational cohorts. This is particularly the case as in this study 8/10 of cases fulfilling McDonald 2010 MS criteria did not meet the criteria from Verhey's predictive model. We have previously shown that 61% of children eventually diagnosed with MS and who had contrast-enhanced scans at ADS onset fulfilled McDonald 2010 MS criteria at first scan.<sup>14</sup> Not all children in this study had spinal imaging and contrast administered, and MRI sequence variation could have affected the sensitivity of MRI.

The potential to diagnose MS at disease onset has implications for MRI protocols for children with suspected ADS, and individual patient treatment decisions. This may be particularly important as new MS drugs are emerging with reported better efficacy in adults than current injectables, and the simultaneous requirement for new drugs to undergo testing in children with the aim of ensuring robust evidence to support authorization by regulatory agencies. To date, eight new MS drugs have been recommended for a Paediatric Investigation Plan,<sup>39</sup> hence earlier diagnosis may increase the eligible clinical trial population. Prognostic and outcome data are needed to inform these decisions. A prospective 2-year follow up of this patient cohort is under way and a further multicentre cohort study with detailed outcome measurements and a biobank have been established.<sup>40</sup> These should yield prognostic information and aid validation of consensus definitions, and the clinical utility of the new McDonald and Verhey proposed MRI criteria for early diagnosis of MS in children.

### Acknowledgments

The funding bodies of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report. MA, CC and EW had full access to all data in the study and had final responsibility for the decision to submit for publication. We also acknowledge the BPSU, supported by the Department of Health (DH), for facilitating the data collection and the reporting clinicians, particularly those who completed the questionnaires. Any views expressed (in publications) are those of the investigator and not necessarily those of the BPSU or DH. We are very grateful to all paediatricians, paediatric neurologists and ophthalmologists who have completed monthly cards, particularly those who have notified cases and returned completed questionnaires and MRI copies to us. We wish to thank all of the secretaries and radiology departments who have assisted with administrative procedures such as posting case report forms and anonymized copies of MRI scans. We are grateful for the laboratory support from Professor Angela Vincent. We also wish to thank Sagar Sedani for help with data collection from the England 2010 index of multiple deprivation database.

### Funding

This study was funded by the Multiple Sclerosis Society UK (grant number 893/08) and Action Medical Research (grant number SP4472). The study also received support from the Wellcome Trust (grant number GN79832) which funded the BCH research facility.

### Conflict of interest statement

MA is funded by the MS Society and Action Medical Research charities. CC has been an investigator on research with unrestricted research funding from Eli Lilly, Wyeth and SHS. WKC has nothing to disclose. CDG has attended the Teva Global MS symposium supported by Teva. KF has nothing to disclose. RG has nothing to disclose. CH has received travel grants from Merck Serono and Bayer. PJ has nothing to disclose. RK has nothing to

disclose. ML has nothing to disclose. MJL has received educational grants to organise a meeting from Novartis, Biogen Idec, Merck Serono and Bayer. KN has nothing to disclose. MGP has received a meeting support grant from Euroimmun. WPW has nothing to disclose. EW has received travel grants from UCB, Shire and Biogen Idec, educational grants to organize meetings from Merck Serono, Novartis, Bayer and Biogen Idec, speaker's fees from Merck Serono and consultancy fees from Genzyme. All authors are members of the UK & Ireland Childhood CNS Inflammatory Demyelination Working Group.

## References

- Orton SM, Herrera BM, Yee IM, et al. Sex ratio of multiple sclerosis in Canada: a longitudinal study. *Lancet Neurol* 2006; 5: 932–936.
- Koch-Henriksen N and Sorensen PS. The changing demographic pattern of multiple sclerosis epidemiology. *Lancet Neurol* 2010; 9: 520–532.
- Renoux C, Vukusic S, Mikaeloff Y, et al. Natural history of multiple sclerosis with childhood onset. *N Engl J Med* 2007; 356: 2603–2613.
- Krupp LB, Banwell B and Tenenbaum S. Consensus definitions proposed for pediatric multiple sclerosis and related disorders. *Neurology* 2007; 68: S7–S12.
- Dean G and Elian M. Age at immigration to England of Asian and Caribbean immigrants and the risk of developing multiple sclerosis. *J Neurol Neurosurg Psychiatry* 1997; 63: 565–568.
- Ebers GC. Environmental factors and multiple sclerosis. *Lancet Neurol* 2008; 7: 268–277.
- Banwell B, Ghezzi A, Bar-Or A, Mikaeloff Y and Tardieu M. Multiple sclerosis in children: clinical diagnosis, therapeutic strategies, and future directions. *Lancet Neurol* 2007; 6: 887–902.
- Dale RC and Pillai SC. Early relapse risk after a first CNS inflammatory demyelination episode: examining international consensus definitions. *Dev Med Child Neurol* 2007; 49: 887–893.
- Mikaeloff Y, Suissa S, Vallee L, et al. First episode of acute CNS inflammatory demyelination in childhood: prognostic factors for multiple sclerosis and disability. *J Pediatr* 2004; 144: 246–252.
- Neuteboom RF, Boon M, Catsman Berrevoets CE, et al. Prognostic factors after a first attack of inflammatory CNS demyelination in children. *Neurology* 2008; 71: 967–973.
- Absoud M, Cummins C, Desai N, et al. Childhood optic neuritis clinical features and outcome. *Arch Dis Child* 2011; 96: 860–862.
- Banwell B, Bar-Or A, Arnold DL, et al. Clinical, environmental, and genetic determinants of multiple sclerosis in children with acute demyelination: a prospective national cohort study. *Lancet Neurol* 2011; 10: 436–445.
- Polman CH, Reingold SC, Banwell B, et al. Diagnostic criteria for multiple sclerosis: 2010 revisions to the McDonald criteria. *Ann Neurol* 2011; 69: 292–302.
- Sedani S, Lim M, Hemingway C, Wassmer E and Absoud M. Paediatric multiple sclerosis: examining utility of the McDonald 2010 criteria. *Mult Scler* 2011; in press.
- Callen DJ, Shroff MM, Branson HM, et al. Role of MRI in the differentiation of ADEM from MS in children. *Neurology* 2009; 72: 968–973.
- Dale RC, de Sousa C, Chong WK, Cox TC, Harding B and Neville BG. Acute disseminated encephalomyelitis, multiphasic disseminated encephalomyelitis and multiple sclerosis in children. *Brain* 2000; 123: 2407–2422.
- Miller DH, Weinshenker BG, Filippi M, et al. Differential diagnosis of suspected multiple sclerosis: a consensus approach. *Mult Scler* 2008; 14: 1157–1174.
- Pohl D, Hennemuth I, von Kries R and Hanefeld F. Paediatric multiple sclerosis and acute disseminated encephalomyelitis in Germany: results of a nationwide survey. *Eur J Pediatrics* 2007; 166: 405–412.
- Banwell B, Kennedy J, Sadovnick D, et al. Incidence of acquired demyelination of the CNS in Canadian children. *Neurology* 2009; 72: 232–239.
- Knowles RL, Smith A, Lynn R and Rahi JS. Using multiple sources to improve and measure case ascertainment in surveillance studies: 20 years of the British Paediatric Surveillance Unit. *J Public Health (Oxf)* 2006; 28: 157–165.
- Poser CM. Revisions to the 2001 McDonald diagnostic criteria. *Ann Neurol* 2006; 59: 727–728.
- Verhey LH, Branson HM, Shroff MM, et al. MRI parameters for prediction of multiple sclerosis diagnosis in children with acute CNS demyelination: a prospective national cohort study. *Lancet Neurol* 2011; in press.
- Callen DJ, Shroff MM, Branson HM, et al. MRI in the diagnosis of pediatric multiple sclerosis. *Neurology* 2009; 72: 961–967.
- Mikaeloff Y, Adamsbaum C, Husson B, et al. MRI prognostic factors for relapse after acute CNS inflammatory demyelination in childhood. *Brain* 2004; 127: 1942–1947.
- Whitworth A. Population Estimates for UK, England and Wales Scotland, and Northern Ireland Mid-2010 Population Estimates. Office for National Statistics, <http://www.ons.gov.uk>, updated 30 June 2011, accessed 11 May 2011.
- Central Statistics Office Ireland. Population estimates for Ireland: regional population projections. Central Statistics Office Ireland <http://www.cso.ie>, updated 4 October 2008, accessed 5 November 2011.
- Whitfield K and Kelly H. Using the two-source capture-recapture method to estimate the incidence of acute flaccid paralysis in Victoria, Australia. *Bull World Health Organ* 2002; 80: 846–851.
- Social Disadvantage Research Centre at the University of Oxford. The English Indices of Deprivation 2010. <http://www.communities.gov.uk/communities/research/indicesdeprivation/deprivation10/>, updated 24 March 2011, accessed 16 November 2011.
- Dean AG SK, Soe MM. OpenEpi: Open Source Epidemiologic Statistics for Public Health, Version 2.3.1. <http://www.OpenEpi.com>, updated 23 June 2011, accessed 11 May 2011.
- Neilson DE, Adams MD, Orr CM, et al. Infection-triggered familial or recurrent cases of acute necrotizing encephalopathy caused by mutations in a component of the nuclear pore, RANBP2. *Am J Hum Genet* 2009; 84: 44–51.
- Langer-Gould A, Zhang JL, Chung J, Yeung Y, Waubant E and Yao J. Incidence of acquired CNS demyelinating syndromes in a multiethnic cohort of children. *Neurology* 2011; 77: 1143–1148.



32. Ghezzi A, Deplano V, Faroni J, et al. Multiple sclerosis in childhood: clinical features of 149 cases. *Mult Scler* 1997; 3: 43–46.
33. Dobson R, Ramagopalan S and Giovannoni G. The effect of gender in clinically isolated syndrome (CIS): a meta-analysis. *Mult Scler* 2011.
34. Torisu H, Kira R, Ishizaki Y, et al. Clinical study of childhood acute disseminated encephalomyelitis, multiple sclerosis, and acute transverse myelitis in Fukuoka Prefecture, Japan. *Brain Dev* 2010; 32: 454–462.
35. Ramagopalan SV, Hoang U, Seagroatt V, et al. Geography of hospital admissions for multiple sclerosis in England and comparison with the geography of hospital admissions for infectious mononucleosis: a descriptive study. *J Neurol Neurosurg Psychiatry* 2011; 82: 682–687.
36. Murray S, Bashir K, Penrice G and Womersley SJ. Epidemiology of multiple sclerosis in Glasgow. *Scott Med J* 2004; 49: 100–104.
37. Absoud M, Parslow RC, Wassmer E, et al. Severe acute disseminated encephalomyelitis: a paediatric intensive care population-based study. *Mult Scler* 2011; 17: 1258–1261.
38. Casrouge A, Zhang SY, Eidenschenk C, et al. Herpes simplex virus encephalitis in human UNC-93B deficiency. *Science* 2006; 314: 308–312.
39. European Medicines Agency. Opinions and decisions on paediatric investigation plans. <http://www.ema.europa.eu>, updated 11 December 2011, accessed 11 December 2011.
40. Absoud M, Cummins C, Chong WK, et al. Paediatric UK demyelinating disease longitudinal study (PUDDLs). *BMC Pediatr* 2011; 11: 68.